

Chagas Disease in Dogs

María Elizabeth Márquez Contreras^{1,*}

¹Laboratorio de Enzimología de Parásitos (LEP), Facultad de Ciencias, Universidad de Los Andes, Mérida, Venezuela

Abstract

Chagas disease is zoonotic illness or an anthrozoosis caused by flagellated protozoan parasite *Trypanosoma cruzi*. This infection presents alarming rates of incidence/prevalence, for this reason, is recognized worldwide as one of the 13 most neglected tropical diseases [1]. Numerous studies have demonstrated the existence of domestic dogs infected with *T. cruzi* across endemic areas ranging from southern United States of America to Argentina [2]. The reported prevalence varies widely (1.42-92%), depending on ecoepidemiological and sociocultural factors [3]. It is important to emphasize that the natural infection in dogs with *T. cruzi* occurs in the same way as in humans, that is to say, through active transmission by vectors, contamination by feces infected with the parasite through wounds or the conjunctiva, can also occur by ingestion of infected vectors or tissues of wild animals present in the peridomicile or home [4]. The transplacental transmission is also an important mode of transmission in dogs [5]. Nevertheless, the main mode of transmission in canine species seems to be the ingestion of infected vectors [6]. During the life cycle of *T. cruzi* the trypomastigotes present in the feces of the triatomines are introduced in the mammalian host by contamination of the insect bite or mucosal membranes. The metacyclic form can penetrate a variety of phagocytic and nonphagocytic nucleated cells. Once inside the cells the parasite becomes in amastigote, which are multiplicative forms that divide into cells. Due to the high parasitic load they produce the lysis of the cells and escapes into the cytoplasm. The amastigotes transform to slender trypomastigotes which can invade adjacent cells, this forms can be ingested by triatomines and they transform into epimastigotes. Finally, after migration to the bug's hindgut, the epimastigotes differentiate into infectious metacyclic trypomastigotes, in this way the life cycle of this microorganism is completed [7].

Corresponding Author : María Elizabeth Márquez Contreras, Laboratorio de Enzimología de Parásitos (LEP), Facultad de Ciencias, Universidad de Los Andes, Mérida, Venezuela, Email: emarquez@ula.ve

Keywords : Anthrozoosis, Ecoepidemiological and sociocultural factors, The transplacental transmission

Received: Oct 16, 2017

Accepted: Jan 3, 2018

Published: Jan 17, 2018

Editor: Mohamed Zowalaty, Virology, Microbiology and Infectious Diseases (VMID) Research Laboratory Bldg E4 Room 505 School of Health Sciences University of KwaZulu-Natal Westville Campus, Durban 4000, Private Bag X54001 South Africa, Email: elzowalaty@ukzn.ac.za

Introduction

Chagas disease is a zoonotic disease caused by flagellated protozoan parasite *T. cruzi*. The infection is recognized worldwide as one of the 13 most neglected tropical diseases [1]. Numerous studies have demonstrated the existence of domestic dogs infected with *T. cruzi* across endemic areas ranging from southern United States of America to Argentina [2]. The reported prevalence varies widely ranging from 1.42 to 92% depending on ecoepidemiological and sociocultural factors [3].

In both humans and dogs the mode of transmission of *T. cruzi* natural infection is through vectors, wound and conjunctiva contamination by feces of contaminated animals, and ingestion of infected vectors or tissues of wild animals living in the peridomicile or home [4]. Transplacental transmission is also an important mode of transmission in dogs [5]. The major mode of transmission in canine species, however, appears to be the ingestion of infected vectors [6].

Life Cycle of *T. cruzi*

In the life cycle of *T. cruzi* the trypomastigotes present in the heces of the triatomines are introduced into the mammalian host by contamination of the insect bite or mucosal membranes. The metacyclic form can penetrate a variety of phagocytic and nonphagocytic nucleated cells. Once inside the cells the parasite becomes an amastigote, which are multiplicative forms that divide into cells. The high parasitic load produces lysis of infected cells and escapes into the cytoplasm. The amastigotes transform to slender trypomastigotes which invade adjacent cells. The trypomastigotes are ingested by triatomines before transforming into epimastigotes, which migrate to the triatomine bug's hindgut and differentiation into infectious forms of metacyclic trypomastigotes completes the life cycle [7].

Clinical Manifestations

Clinical manifestations in dogs range from asymptomatic, acute myocarditis, chronic progressive cardiac disease to sudden death [8, 3]. The clinical signs such as the development of diffuse chronic myocarditis with histological and electrocardiographic changes in Chagas-infected dogs during both the chronic and acute stages closely resemble the symptoms of human

disease [4]. This is the reason why canine species are considered the experimental model of choice for human Chagas disease study [2]. Acute illness has been reported more frequently in very young dogs (less than 1 year old) and generally involves myocarditis and cardiac arrhythmias [3]. The chronic phase of Chagas' disease in canine species chagasic, can cause severe cardiac damage involving dilated cardiomyopathy, electrocardiogram (ECG) abnormalities, and cardiac failure [9]. This clinical presentation is highly variable and depends on the type of strain, route of infection, and parasite burden [10].

Diagnosis of Chagas Disease Canine

The diagnosis is based on serology [11], direct or indirect parasitological tests, and molecular tests [12]. Parasitological diagnosis is made by microscopic examination of either the lymph node aspirates of blood, or cerebrospinal fluid (CSF) of infected dogs [13]. Polymerase chain reaction (PCR) technique could be used on any patient sample that contains trypanosome DNA [14].

Epidemiology of Chagas Disease in Canines

Canine species are considered an important reservoir of *T. cruzi* in the domestic cycle carrying a risk for human population; because, these animals living in close proximity to humans are within the transmission and maintenance cycles of the parasite and serve as a source of food for insect vectors [2,15]. Reservoirs may be considered as a complex ecological system consisting of one or more species responsible for maintaining a given parasite species in nature. The infestation of houses and peridomestic areas is considered to be a major risk factor for Chagas disease transmission. Dog infection with *T. cruzi* is an important veterinary health concern in many countries of South America and more recently in the United States, in which the seroprevalence of specific anti-*T. cruzi* antibodies has also been reported [16, 17]. Infected vectors have been reported in Texas military kennels, where some dogs showed clinical signs compatible with Chagas disease infection [18]. From the veterinary point of view, it is also important to consider that dogs are susceptible to acquire American trypanosomiasis characterized by heart conditions, such as electrical conduction disturbances, and ventricular and supraventricular arrhythmias, as well as secondary signs such as ascites,

respiratory distress, thoracic effusion, and cyanosis. Texas is a high-risk state for transmission of the parasite to dogs, considering the diversity of triatomine vectors, reservoir hosts, and previous documentation of canine disease [19]. A high seroprevalence of canine infection with *T. cruzi* has been detected from southeastern USA States of Louisiana, Oklahoma, Georgia, and Texas; showing the epidemiological role of the domestic dog [20]. In canines from countries in Latin American, various reports showed seroprevalences from 24% to 65% [21, 22]. The presence of infected canine species increases the risk of transmission of parasites to vectors, and thus the probability of infection in humans. Generally, canine infections are more prevalent than human infections, a fact likely related to oral *T. cruzi* transmission, a more efficient route of infection and apparently common to many mammals [23], and also the fact that dogs often sleep near houses and may come in greater contact with peridomestic vectors.

It has been considered that separating domestic animals, mainly dogs, from people can significantly reduce transmission to humans [24]. The fact that there is a high prevalence of canine infections represents a warning signal that should lead to the planning and execution of effective vector control strategies, i.e. it is essential to carry out fumigation strategies of areas infested with triatomines, to exterminate them and thus eliminate transmission [25].

In studies carried out by Meyer et al. (2017) [26] in along border Texas-México, the dogs showed an overall apparent seroprevalence of 7.4-18.9%. This values are similars to that described in dogs of endemic populations for Chagas in Peru (12.3%) [27], Argentina (45.6%) [11], Panama (11.1%) [28], Costa Rica (27.7%) [29], Yucatan State, Mexico (9.8%-14.4%) [30] and Mexico State, Mexico (10%-15.8%) [31].

The dogs are targeted in a Chagas' disease control strategy, because they are considered natural sentinels [26]. These animals generally inhabit a defined territory; are accessible, easy to enumerate and capture, and their population allows representative sampling [31].

Control Measurement

Control measures to prevent transmission of Chagas' disease in endemic areas include: removing litter from the surroundings of the house, keeping clean

areas where dogs stayed longer, and conduct periodic assessments to ensure the animals are free of insects. Placing meshes on doors and windows, and covering cracks in the whole house helps prevent the entry of triatomines to the home and minimizes their contact with humans or canine species.

References

1. Hotez P, Molyneux D, Fenwick A, Kumaresan J, Sachs S, Sachs J, Savioli L (2007) Control of neglected tropical diseases. *N Engl J Med* 357 (10):1018-1027
2. Gurtler R, Cecere M, Lauricella M, Cardinal M, Kitron U, Cohen J. (2007). Domestic dogs and cats as source of *Trypanosoma cruzi* infection in rural northwestern Argentina. *Parasitology* 134:69-82
3. Kjos S, Snowden K, Craig T, Lewis B, Ronald N, Olson J (2008) Distribution and characterization of canine Chagas disease in Texas. *Vet Parasitol* 152 (3-4):249-256
4. Barr S (2009) Canine Chagas' disease (American Trypanosomiasis) in North America. *Vet Clin North Am Small Anim Pract* 39(6):1055-1064
5. Castañera M, Lauricella M, Chuit R, Gürtler R (1998) Evaluation of dogs as sentinels of the transmission of *Trypanosoma cruzi* in a rural area of north-western Argentina. *Ann Trop Med Parasitol* 92 (6):671-683
6. Eloy L, Lucheis S (2009) Canine trypanosomiasis: etiology of infection and implications for public health. *J Venom Anim Toxins Incl Trop Dis* 15(4): 589-611
7. Tyler K, Engman D (2001) The life cycle of *Trypanosoma cruzi* revisited. *Int J Parasitol* 31(5-6):472-481
8. Barr S (2009) Canine Chagas' disease (American trypanosomiasis) in North America. *Vet Clin North Am Small Anim Pract* 39(6):1055-1064
9. Barr S, Simpson R, Schmidt S, Bunge M, Authement J, Lozano F (1989) Chronic dilatative myocarditis caused by *Trypanosoma cruzi* in two dogs. *J Am Vet Med Assoc* 195(9):1237-1241
10. González-Vieyra S, Ramírez-Durán N, Sandoval-Trujillo Á, Vázquez-Chagoyán J, Monroy-Salazar H, Barbabosa-Pliego A (2011) *Trypanosoma cruzi* in dogs: electrocardiographic and

- echocardiographic evaluation, in Malinalco, State of Mexico. *Res Rep Trop Med* 2:155-161
11. Enriquez G, Cardinal M, Orozco M, Schijman A, Gürtler R (2013) Detection of *Trypanosoma cruzi* infection in naturally-infected dogs and cats using serological, parasitological and molecular methods. *Acta Trop* 126(3): 211-217
 12. Lizundia R, Picado A, Cordero M, Calderón A, Deborggraeve S, Montenegro V, Urbina A (2014) Molecular and serological rapid tests as markers of *Trypanosoma cruzi* infection in dogs in Costa Rica *Trop Parasitol* 4(2):111-114
 13. Fujita O, Sanabria L, Inchausti A, De Arias A, Tomizawa Y, Oku Y (1994) Animal reservoirs for *Trypanosoma cruzi* infection in an endemic area in Paraguay. *J Vet Med Sci* 56(2):305-308
 14. Araújo F, Bahia M, Magalhães N, Martins-Filho O, Veloso V, Carneiro C, Tafuri W, Lana M (2002) Follow-up of experimental chronic Chagas' disease in dogs: use of polymerase chain reaction (PCR) compared with parasitological and serological methods. *Acta Trop* 81(1):21-31
 15. Ortega Pacheco A, Guzmán Marín E, Acosta Viana K, Vado Solís I, Jiménez Delgadillo B, Cárdenas Marrufo M, Pérez-Osorio C, Puerto-Solís M, Jiménez-Coello M (2017) Serological survey of *Leptospira interrogans*, *Toxoplasma gondii* and *Trypanosoma cruzi* in free roaming domestic dogs and cats from a marginated rural area of Yucatan Mexico. *Vet Med Sci* 3(1):40-47
 16. Villalobos G, Martínez-Hernández F, de la Torre P, Lactette J, Espinoza B (2011) Entomological indices, feeding sources, and molecular identification of *Triatoma phyllosoma* (Hemiptera: Reduviidae) one of the main vectors of Chagas disease in the Istmo de Tehuantepec, Oaxaca, Mexico. *Am J Trop Med Hyg* 85(3):490-497
 17. Bern C, Kjos S, Yabsley M, Montgomery S (2011) *Trypanosoma cruzi* and Chagas' Disease in the United States. *Clinical Microbiology Reviews*. 24 (4):655-681
 18. Arce-Fonseca M, Carrillo-Sánchez, S, Molina-Barrios R, Martínez-Cruz M, Cedillo-Cobián J, Henao-Díaz A, Rodríguez-Morales O (2017) Seropositivity for *Trypanosoma cruzi* in domestic dogs from Sonora, Mexico. *Infect Dis Poverty* 6(1):120
 19. Tenney T, Curtis-Robles R, Snowden K, Hamer S (2014) Shelter dogs as sentinels for *Trypanosoma cruzi* transmission across Texas. *Emerg Infect Dis* 20 (8):1323-1326
 20. Mc Phatter L, Roachell W, Mahmood F, Hoffman L, Lockwood N, Osuna A, Lopez J, Debboun M Haro-Rodriguez S, Ventura-Rodriguez LV, Noguera-Torres B, Bustos-Saldaña R (2012) Vector surveillance to determine species composition and occurrence of *Trypanosoma cruzi* at three military installations in San Antonio, Texas. *US Army Med Dep J* 3:12-21
 21. Kjos S, Snowden K, Olson J (2009) Biogeography and *Trypanosoma cruzi* infection prevalence of Chagas disease vectors in Texas, USA. *Vector Borne Zoonotic Dis* 9(1):41-50
 22. Curtis-Robles R, Snowden K, Dominguez B, Dinges L, Rodgers S, Mays G, Hamer S (2017) Epidemiology and molecular typing of *Trypanosoma cruzi* in naturally-infected hound dogs and associated triatomine vectors in Texas, USA. *Plos Negl Trop Dis* 11(1):e0005298
 23. Beard C, Pye G, Steurer F, Rodriguez R, Campman R, Peterson A, Ramsey J, Wirtz R, Robinson L (2003) Chagas disease in a domestic transmission cycle, southern Texas, USA. *Emerg Infect Dis* 9 (1):103-105
 24. Estrada-Franco, J, Bhatia V, Diaz-Albiter H, Ochoa-Garcia L, Barbabosa A, Vazquez-Chagoyan J, Martinez-Pérez M, Guzman-Bracho C, Garg N (2006) Human *Trypanosoma cruzi* infection and seropositivity in dogs, Mexico. *Emerg Infect Dis* 12(4):624-630
 25. Barbabosa-Pliego A, Gil P, Hernández D, Aparicio-Burgos J, de Oca-Jiménez R, Martínez-Castañeda J, Ochoa-García L, Guzmán-Bracho C, Estrada-Franco J, Garg N, Chagoyán J (2011) Prevalence of *Trypanosoma cruzi* in dogs (*Canis familiaris*) and triatomines during 2008 in a sanitary region of the state of Mexico, Mexico. *Vector Borne Zoonotic Dis* 11(2):151-156
 26. Meyers A, Meinders M, Hamer S (2017) Widespread *Trypanosoma cruzi* infection in government working dogs along the Texas-Mexico border: Discordant serology, parasite genotyping and associated vectors. *PLoS Negl Trop Dis* 11(8):e0005819

27. Castillo-Neyra R, Chou Chu L, Quispe-Machaca V, Ancca-Juarez J, Malaga Chavez F, Bastos Mazuelos M, Naquira C, Bern C, Gilman R, Levy M (2015) The potential of canine sentinels for reemerging *Trypanosoma cruzi* transmission. *Prev Vet Med* 120 (3-4):349-356
28. Pineda V, Saldaña A, Monfante I, Santamaría A, Gottdenker N, Yabsley M, Rapoport G, Calzada J (2011) Prevalence of trypanosome infections in dogs from Chagas disease endemic regions in Panama, Central America. *Vet Parasitol* 178(3-4):360-363
29. Montenegro V, Jimenez M, Dias J, Zeledon R (2002) Chagas disease in dogs from endemic areas of Costa Rica. *Mem Inst Oswaldo Cruz* 97(4):491-494
30. Jimenez-Coello M, Poot-Cob M, Ortega-Pacheco A, Guzmán-Marín E, Ramos-Ligonio A, Sauri-Arceo C, Acosta-Viana K (2008) American trypanosomiasis in dogs from an urban and rural area of Yucatan, Mexico. *Vector Borne Zoonotic Dis* 8(6):755-761
31. Mejía A, Portugal-García C, Chávez-López V, García-Vázquez Z, Ramos C (2017) Evidencia serológica de infección por *Trypanosoma cruzi* en perros atendidos en clínicas veterinarias del área conurbada de Cuernavaca, Morelos. *Salud Públ Méx* 59(3):205-206