

Lipopolysaccharide Prompts Oxidative Stress and Apoptosis in Rats' Testicular Tissue

Amal A Halawa^{1*}, Mohamed A El-Adl², Mohamed F Hamed³, Ahmed Z Balboula⁴, Mohammed A Elmetwally⁴

¹Department of Forensic Medicine and Toxicology, Faculty of Veterinary Medicine, Mansoura University, Mansoura, Egypt. am_abdelmonem@yahoo.com

²Department of Biochemistry and Chemistry of Nutrition, Faculty of Veterinary Medicine, Mansoura University, Mansoura, Egypt.

³Department of Pathology, Faculty of Veterinary Medicine, Mansoura University, Mansoura, Egypt.

⁴Department of Theriogenology, Faculty of Veterinary Medicine, Mansoura University, Mansoura, Egypt.

Abstract

Lipopolysaccharide (LPS) is a component of the outer membrane of gram negative bacteria. LPS challenging allows switching transcription of proinflammatory cytokines on via over stimulation of Toll-like receptors (TLRs) signaling pathway with subsequent pathogenic inflammatory response. We investigated the possible reproductive toxicity of LPS in male Wister albino rats. Oxidative stress markers, antioxidant status and caspase-3 activity were analyzed in testicular tissues of rats exposed to either saline or LPS (4 mg/kg BW, ip; 0.18 of the LD₅₀). The samples were collected at 6 h and 72 h after injection of LPS. A significant reduction in testicular reduced glutathione (GSH), glutathione-S-transferase (GST) and superoxide dismutase (SOD) was observed at 72 h compared to control group. Total antioxidant capacity was decreased at 6 h with additional significant reduction at 72 h. Catalase activity was reduced significantly at both 6 and 72 h. Malondialdehyde (MDA) was increased ($P \leq 0.05$) in LPS injected rats without variation between 6 and 72 h. A significant increase in nitric oxide (NO) was observed at 72 h after injection. A time-dependent increase in LPS-treated groups was observed in the concentration of caspase-3. Histopathological analysis revealed degenerative changes and necrosis of seminiferous tubules after 6 h with further accumulation of eosinophilic edematous transudate in its lumen after 72 h. In conclusion, by increasing time of exposure, LPS induced lipid peroxidation, oxidative stress, reduced testicular antioxidant capacity and encouraged testicular apoptosis which could be possible mechanisms for impairment of testicular function.

Corresponding author: Amal A Halawa, Department of Forensic Medicine and Toxicology, Faculty of Veterinary Medicine, Mansoura University, 35516 Mansoura, 60 Elgomhoria St, Egypt. Tel: 01020992886, Fax: +20502200696. Email: am_abdelmonem@yahoo.com

Running title: LPS adversely affects rat testicular tissue

Keywords: Lipopolysaccharide, Oxidative stress, Caspase 3, Reproductive toxicity, Testis

Received: Feb 28, 2018

Accepted: Mar 22, 2018

Published: Mar 29, 2018

Editor: BURAK DiK, Selcuk University, Veterinary Faculty, Farmacology And Toxicology, TURKEY.

Introduction

Lipopolysaccharide (LPS) is a glycolipid component of the cell wall of gram negative bacteria such as *Escherichia* and *Salmonella* species. The name of endotoxin is due to the biological activities that induced by LPS after entering the host organism [1]. LPS consists of lipid A part which is responsible for the toxic proinflammatory properties of LPS [2,3] and polysaccharide side chain part, which consists of a core oligosaccharide and the terminal O-specific chain antigen [4,5] that determines the serological specificity or the bacterial serotype [6]. LPS is considered as a molecular pattern related to the pathogen-associated molecular patterns (PAMPs) via which the bacteria can be recognized by specific host receptors called pattern recognition receptors (PRRs) [7]. LPS performs as a toxin via over stimulation of Toll-like receptors (TLRs) signaling pathway, that promotes nuclear translocation of NF- κ B [8] and switches on transcription of proinflammatory cytokines such as IL-1 β , IL-6 and TNF α that encourages pathogenic inflammatory responses [5, 9]. Exposure to high doses of LPS triggers the production of the proinflammatory mediators which may result in a harmful condition termed oxidative stress [10, 11]. Reactive oxygen species (ROS) are thought to be involved in the mechanism of LPS toxicity [12].

Male infertility is essentially a result of disruption in spermatogenesis process, which is based on reduction in sperm count, motility and viability, oxidative stress and disturbance in sex hormones [13]. Infections lead to an extreme production of ROS, resulting in an oxidative stress which impairs the sperm functions, as well as fertilization [14]. It was suggested that LPS inhibits human sperm motility by decreasing intracellular cAMP [15]. Previous study evaluated the adverse effects of LPS on male reproduction system identifying increased oxidative stress and proinflammatory stress markers in the testes leading to marked testicular dysfunction [16].

We aimed in the present study to estimate the potential mechanisms for LPS to induce testicular dysfunction via measuring the antioxidant activity and oxidative stress, in addition the effect of LPS on the apoptotic gene marker, Caspase-3 in male albino rats. Histopathological analysis on testicular structure was also applied.

Materials and Methods

Chemicals

Lipopolysaccharide (LPS) (*E. coli*, serotype 0111:B4; Sigma-Aldrich) was dissolved in distilled water. Rats in control group were administered intraperitoneal pyrogen-free 0.9% NaCl.

Animals

Fifteen male Wister albino rats (3 months old) were purchased from Faculty of Medicine, Mansoura University, Mansoura, Egypt. The animals were allowed for accommodation at 2 weeks for the laboratory conditions (12 h light/dark cycles) before commencement of the experiment. Animals were received feed and water *ad libitum*. Animals received human care in compliance with the guidelines of animal care of the National Institutes of Health, and all animals applications were performed in accordance with the Ethics Committee of the National Research Centre, Egypt, registration number (09/189).

Experimental Design

A total of 15 rats were divided into control (n = 5) and LPS treated groups (n = 10). In the second group, testes were collected after 6 h (n=5) and after 72 h (n=5). Control rats received intraperitoneal injection of pyrogen-free 0.9% NaCl. According to Nezić et al (2009) [17], single intraperitoneal injection of non-lethal dose of LPS was administered to the group treated with LPS (4 mg/kg BW; 0.18 of the LD₅₀ [22.15 mg/kg]) [17] Rats were sacrificed and testes were collected at 6 and 72 h time points after LPS injection.

Tissue Preparation

One g of each testis was washed by PBS (pH 7.4) and then homogenized in nine volume of PBS. Tissue homogenate was further centrifuged at 4000 rpm for 20 minutes at 4°C and stored at -80°C according to Fernandez-botran et al. (2002) [18] till antioxidant and oxidative stress markers were performed.

Biochemical Analysis:

Testicular Reduced Glutathione Concentration:

Reduced glutathione in testicular homogenate was determined using the method of Beutler et al. (1963) [19]. 5, 5 dithiobis 2- nitrobenzene (DTNB) was

used to give yellow colored complex in conjugation with reduced glutathione

Testicular Glutathione-S-tTransferase Activity:

GST activity was determined by measuring the conjugation of 1- chloro- 2, 4-dinitrobenzene (CDNB) with reduced glutathione according to method of Habig et al. (1976) [20].

Testicular Superoxide Dismutase Activity:

Tissue homogenate of testis was mixed with working solution contained with 0.5mM phosphate buffer, 0.3mM NADPH+ and 0.5mM nitroblue tetrazolium for a min then the reaction was further inhibited by phenazine methosulphate. The percent of inhibition of superoxide radical was estimated and multiplied by 3.75 to determine SOD activity according to Nishikimi et al. (1972) [21].

Total Antioxidant Capacity:

Capacity of total antioxidant capacity was determined according to Koracevic et al. (2001) [22]. Capacity of total antioxidant was determined with conjugation of hydrogen peroxide with 3,5,dichloro-2-hydroxy benzensulphonate to form pinky color compound that inversely proportional with total antioxidant capacity of samples.

Testicular Catalase Activity:

According to Aebi (1984) [23], catalase activity was determined through the enzymatic reaction initiated via binding of the produced hydrogen peroxide with 3,5-Dichloro-2-hydroxybenzene sulphonic acid and 4-aminophenazone. The produced colored complex is inversely proportional with catalase activity in tissues.

Oxidative Stress Marker (Lipid peroxidation):

The levels of lipid peroxidation expressed as malondialdehyde concentration was measured spectrophotometrically followed the method of Draper and Hadley (1990) [24]. Thiobarbituric acid reacts with MDA in acid medium to form reactive product. The absorbance of the resultant pink product can be measured at 534 nm.

Nitrosative Stress Marker:

The level of nitrite was used to determine the extent of nitrosative stress on testicular tissues according to the method of Montgomery and Dymock

(1961) [25]. In the presence of nitrite and in an acidic medium, the formed nitrous acid diazotized sulphanilamide and the product is coupled with N-(1-naphthyl) ethylenediamine. The resulting azodye has a bright reddish-purple color which can be measured at 540 nm.

Flowcytometric Determination of Caspase 3:

The determination of caspase as an apoptotic marker was occurred using flow cytometric technique according to technique of Tribukait (1987) [26]. Testicular tissues were washed in isotonic tris EDTA buffer (3.029 g of 0.1 M tris (hydroxymethyl aminomethane , 1.022 g of 0.07 M sodium chloride and 0.47 g of 0.005 M EDTA) and were further suspended in absolute ethanol for accomplishing of fixation step of testicular tissues. The fixed tissues was conjugated by adequate dilution (1:50) of caspase-3 primary antibody (Abcam, USA), that was further mixed with secondary antibodies (IgG) (Abcam, USA) with the appropriate dilution (1:2000). Finally, cells were suspended in solution containing PBS/BSA. After staining procedures were occurred, measurements of positive cells was determined using flow cytometer device (FACS caliber flow cytometer "Becton Dickinson , Sunnyvale , CA, USA) equipped with a compact air cooled low power 15 mwatt argon iron laser beam (488 nm).

Histopathological Examination:

The testes were dehydrated in graded ethanol and embedded in paraffin wax. 5 µm thick paraffin embedded sections were cut and routinely stained with hematoxyline and eosin according to Bancroft et al (1996) [27]. Each section was examined by light microscopy.

Statistical Analysis:

The results were expressed using one way ANOVA to determine the difference between means in every group. Least significance difference using Duncan's post-hoc test was used to determine statistically significance difference between means at a significant level 0.05 where $P \leq 0.05$. All data were reported as means \pm SEM. Statistical analyses were done using SAS® (version 9.2, SAS Institute, Cary, NC, USA). For all analyses, $P \leq 0.05$ was defined as significant.

Results

Effects of LPS on Antioxidant Activity

Antioxidant Status:

Glutathione concentration (GSH) showed a significant reduction at 72 h (15.42 ± 0.678 mg/g) after LPS injection compared to 6 h (17.58 ± 0.174 mg/g) exposure and control (18.17 ± 0.593 mg/g) groups (Figure 1a). GST showed a significant decrease after 72 h (0.677 ± 0.009 U/g) of LPS exposure in comparison with control (1.343 ± 0.015 U/g) and 6 h (0.967 ± 0.152 U/g) exposure ($P \leq 0.05$) (Figure 1b). SOD activity was reduced at 6 h (238.82 ± 7.78 U/g) without significant variation compared to control (242.13 ± 7.83 U/g) while the reduction at 72 h (207.81 ± 8.14 U/g) was significantly varied compared to control and 6 h groups ($P \leq 0.05$) (Figure 1c).

Total Testicular Antioxidant Capacity:

A significant reduction in total antioxidant capacity was observed at 6 h (0.66 ± 0.017 μ mol/g) with additional decline at 72 h (0.203 ± 0.026 μ mol/g) compared to control group (0.947 ± 0.009 μ mol/g) ($P \leq 0.05$) (Figure 2a). Corresponding to antioxidant capacity of rats exposed to LPS, the activity of catalase was decreased significantly at 6 h (0.187 ± 0.035 U/g) and 72 h (0.17 ± 0.012 U/g) compared to control group (0.317 ± 0.019 U/g) ($P \leq 0.05$) without significant variation between LPS-treated groups (Figure 2b).

Oxidative Stress Marker:

The levels of lipid peroxidation (MDA) in rats exposed to LPS showed a significant increase after 6 h (12.41 ± 0.49 nmol/g) and 72 h (10.62 ± 0.50 nmol/g) of exposure in comparison with control group (7.38 ± 0.50 nmol/g) ($P \leq 0.05$). Although there was a reduction in testicular MDA levels after 72 h compared to 6 h, but this reduction was non-significant ($P > 0.05$) (Figure 3a).

Nitrosative Stress Marker:

A robust increase occurred ($P \leq 0.05$) in testicular nitrite levels after 72 h (35.4 ± 1.12 nmol/g) after LPS injection compared to control group (25.61 ± 3.45 nmol/g). The increase of nitrite level at 6 h was insignificant (31.2 ± 0.46 nmol/g) (Figure 3b).

Effect of LPS on Apoptotic Factor Caspase -3

The concentration of caspase-3 showed a marked increase ($P \leq 0.05$) at either 6 h (32.24 ± 2.15) or 72 h (54.74 ± 3.29) compared to control group (15.12 ± 1.22) (Figure 4).

Histopathological Analysis of Testicular Tissue

Seminiferous tubules showed normal spermatogenic epithelium, Sertoli cells and interstitial tissue in control group (Figure 5a), whereas testes collected 6 h after LPS injection displayed degenerative changes and necrosis of seminiferous tubules (Figure 5b). After 72 h, necrosis of spermatogenic epithelium lining seminiferous tubules and accumulation of eosinophilic edematous transudate in its lumen were observed (Figure 5c).

Discussion

Testicular tissues may lack an adequate oxygen supply, so testes may suffer dramatically from oxidative stress due to the high levels of unsaturated fatty acids and excessive production of ROS that could be produced from leakage of mitochondrial membrane and might impact both spermatogenic and steroidogenic functions [28, 29]. At the testicular level, oxidative stress can be modulating steroidogenic capacity of Leydig cells [30] besides disrupting ability of germinal epithelium to differentiate spermatozoa [31] which may result in infertility [32].

The capability of LPS for induction of oxidative stress in testicular tissues was previously studied in mice [33]. It was reported that the LPS could possibly induce oxidative stress in rats due to the disruption in germ cell layer in seminiferous tubules which increased the production of proinflammatory cytokines and generation of ROS [34]. Moreover, the LPS would possibly increase the activity and concentration of nitric oxide synthetase (NOS) in testes that produced excessive amount of nitric oxide causing nitrosative stress in adult rats [35]. We aimed to reveal the ability of LPS to induce oxidative stress as well as apoptosis in short periods of time (6 and 72 h) in testicular tissue of rats as possible mechanisms for testicular dysfunction.

In the current study, GSH concentration showed a significant decrease after 72 hours of LPS treatment, and occurred a decrease in GST activity in comparison with control group. It was shown the induced infection




Figure 1: a) GSH concentration, b) GST activity and c) SOD activity in control and LPS-treated groups (4 mg/kg BW ip). Results are expressed as mean \pm SEM. The different letters are statistically significant ($P \leq 0.05$).




Figure 2: a) Total antioxidant capacity and b) Catalase activity in control and LPS-treated groups (4 mg/kg BW ip). Results are expressed as mean \pm SEM. The different letters are statistically significant ($P \leq 0.05$).




Figure 3: a) MDA concentration and b) NO concentration in control and LPS-treated groups (4 mg/kg BW ip). Results are expressed as mean \pm SEM. The different letters are statistically significant ($P \leq 0.05$).




Figure 4: Relative expression of caspase-3 in testicular tissues of either control or LPS-treated rats (4 mg/kg BW ip). Results are expressed as mean \pm SEM. The different letters are statistically significant ($P \leq 0.05$).




Figure 5: Histopathological results. a) Control group: seminiferous tubules with normal spermatogenic epithelium (arrow), normal Sertoli cells and normal interstitial tissue. b) After 6 hours seminiferous tubules display degenerative changes and necrosis of spermatogenic epithelium (arrow) and abnormal Sertoli cells. c) After 72 h necrosis of spermatogenic epithelium lining seminiferous tubules and accumulation of eosinophilic edematous transudate in its lumen.

using LPS was capable of encouraging oxidative stress that concurrently would result in decrease GSH in human dendritic cells [36]. In another study, GST activity showed a significant decrease after *Cyanobacterium* LPS treatment in zebra fish [37]. Similarly, the activity of hepatic GST showed a significant decline when galactosamine /lipopolysaccharide were injected to rats [38].

As a result of the oxidative damage and generation of free radicals, the concentration of total antioxidant capacity revealed a significant gradually reduction by the increase with time of LPS exposure. In a similar study, rats administered 30 mg/kg LPS showed a significant reduction in total antioxidant capacity that was measured as ferric reducing ability of plasma [39]. The reduction of antioxidant capacity was followed by a significant decrease in testicular catalase activity by 54% after 72 hours of treatment which was further confirmed by the study of Spolarics and Wu (1997) [40], whom found that LPS treatment would result in a significant reduction in catalase activity by 45% in Kupffer cells [40]. Such reduction reflects the inability of testicular cells to eliminate H₂O₂ produced by the inflammatory events resulting from LPS injection or may be due to enzyme activation caused by ROS generated in testicular cells [41, 42].

SOD activity showed a significant decrease after 72 hours of LPS treatment which was observed in broilers treated with 0.5 mg/ml LPS attributing to disturbance in antioxidant defense system [43].

One of the various effects of oxidative stress is peroxidation of unsaturated fatty acids in membrane. The degree of lipid peroxidation, measured as MDA, was used to explain the role of LPS in peroxidation of lipids in testicular tissues. The concentration of MDA was increased immediately after 6 h of LPS administration with slight non-significant reduction at 72 h. Testicular MDA levels increased at the same dose of LPS at 6 h and returned to normal level at 12 and 72 h [44]. Likewise, 5 mg/kg LPS increased MDA levels in albino rats with reduction in testicular antioxidant activities [41]. A robust increase the levels of lipid peroxidation were observed with higher doses of LPS in Fischer rats [45]. Also, LPS (200 µg/mouse, for 2 h) induced lipid peroxidation products (MDA and 4-hydroxynonenal) in

isolated mice Leydig cell homogenates [46]. Moreover, LPS- treated Sertoli cells of adult rats (50 µg/ml for 12 h) showed augmentation in H₂O₂ and lipid peroxidation with reduction in activities and concentration of the antioxidant parameters (GSH, GST, SOD, CAT) [47]. Ability of LPS (1 µg/ml, for 12 h) to exert oxidative stress was also verified using primary culture of rat Leydig cells [48].

The levels of nitrosative stress were determined using the levels of nitric oxide production in testicular tissues. NO was increased significantly after 72 h of treatment which was observed as well in macrophage cell line due to induction of nitric oxide synthetase expression in macrophage [49]. In a study performed by O'Bryan et al. (2000) [50], testicular tissues of rats showed a significant increase in NO production due to up-regulation of inducible nitric oxide synthetase (iNOS) after LPS administration [50].

Stangelet al. (1996) [51] found that ROS and NO were capable of inducing apoptosis in rat skeletal myoblast [51]. Apoptosis is a programmed cell death which occurs physiologically in testicular germ cells to balance their number to that of Sertoli cells [52]. Apoptosis is generated by various pathways, all initiate Caspase pathway (Caspase 8 and 9) that in turn activate the effector Caspase 3 [53, 54]. The results of the present study showed early induction of apoptosis through increase in Caspase-3 concentration (6 h), which could be due to the induction of testicular oxidative stress as well as increase in NO levels. Pro-inflammatory cytokines that were induced by LPS is another mechanism by which apoptosis could be induced through TNF-dependent pathway [55]. Besides, LPS was capable of inducing apoptosis in hepatocyte which was followed by an increase in production of Caspase-3 in co-culture of hepatocyte and Kupffer cells [56]. LPS administered to mice (intraperitoneal, 0.1 mg/kg, 7 days) caused increase of apoptotic germ cells via Fas/ FasL system up to 5 weeks after LPS exposure and observed peak at 24 h [57]. Additionally, Leydig cells incubated for 12 h with LPS was showed increase in Caspase 3 activity [58].

The aforementioned alterations that were observed in the present work were supported by the histopathological examination of the testicular tissue.

After 6 h, seminiferous tubules showed necrosis and degenerative changes, meanwhile, testes collected 72 h post LPS injection, showed necrosis of spermatogenic epithelium lining seminiferous tubules with eosinophilic edematous transudate in its lumen. Similarly, at the same dose of LPS, accumulation of immature germ cells in the lumen of seminiferous tubules was observed after 6 h of LPS injection. While at 72 h, large numbers of immature germ cells were observed in the lumen of epididymis with intercellular gaps in seminiferous epithelium [44]. The histopathological lesions in rat testes after single intraperitoneal injection of LPS revealed normal structure of seminiferous tubules at 3 and 6 h after LPS injection; whereas seminiferous epithelium showed degeneration and sloughing at 24 and 72 h time points [41]. At the same dose of LPS, seminiferous tubules showed thinning in its epithelium lining disordered architecture and uneven arrangement of spermatogenic cells, together with reduced number of mature sperms and noticeable shedding of spermatogenic cells in the lumens of seminiferous tubules were observed. Such lesions become more pronounced by 72 h after LPS injection [59].

The data of the current work are in consistence with other studies proved that oxidative stress is considered as a key factor in steroidogenesis and spermatogenesis alteration with subsequent infertility [41, 50, 60].

In conclusion LPS can induce oxidative stress after short period from exposure as early as 6 h and continue up to 72 h resulting from a significant decrease in antioxidant defense enzymes with an increase in lipid peroxidation and NO levels. The LPS-induced oxidative stress could be accompanied with apoptosis through activation of Caspase-3 resulting in impairment in testicular functions. This effect is a crucial for the reproductive performance and future fertility of male due to the LPS effect on germinal epithelium of testes.

Acknowledgements

The authors appreciate everyone participate in the present work.

Funding

The author(s) received no financial support.

Conflict of interest

The authors declare no conflict of interest.

References:

1. Kabanov, D.S., Prokhorenko, I.R., 2010. Structural analysis of lipopolysaccharides from Gram-negative bacteria. *Biochem.* 75, 383-404.
2. Raetz, C.R.H., Garrett, T.A., Reynolds, C.M., Shaw, W.A., Moore, J.D., et al., 2006. Kdo2-Lipid A of *Escherichia coli*, a defined endotoxin that activates macrophages via TLR-4. *J. Lipid Res.* 47, 1097-1111.
3. Kirschning, C.J., Bauer, S., 2001. Toll-like receptors: cellular signal transducers for exogenous molecular patterns causing immune responses. *Int. J. Med. Microbiol.* 291, 251-260.
4. Hodgson, J.C., 2006. Endotoxin and mammalian host responses during experimental disease. *J. Comp. Pathol.* 135, 157-175.
5. Tomlinson, J.E., Blikslager, A.T., 2004. Interactions between lipopolysaccharide and the intestinal epithelium. *J. Am. Vet. Med. Assoc.* 224, 1446-1452.
6. Alexander, C., Rietschel, E., 2001. J. Bacterial lipopolysaccharides and innate immunity. *Endotoxin. Res.* 7, 167-202.
7. Eckmann, L., 2004. Innate immunity and mucosal bacterial interactions in the intestine. *Curr. Opin. Gastroenterol.* 20, 82-88.
8. Brightbill, H.D., Modlin, R.L., 2000. Toll-like receptors: molecular mechanisms of the mammalian immune response. *Immunology.* 101, 1-10.
9. Gutschmann, T., Schromm, A.B. and Brandenburg, K., 2007. The physicochemistry of endotoxins in relation to bioactivity. *Int. J. Med. Microbiol.* 297, 341-352.
10. Libby, P., 2007. Inflammatory mechanisms: the molecular basis of inflammation and disease. *Nutr. Rev.* 65, S140-S146.
11. Victor, V.M., Rocha, M. and De la Fuente, M., 2004. Immune cells: free radicals and antioxidants in sepsis. *Int. Immunopharmacol.* 4, 327-347.
12. Kallapura, G., Pumford, N.R., Hernandez-Velasco, X. Hargis, B.M., Tellez, G., 2014. Mechanisms involved in lipopolysaccharide derived ROS and RNS oxidative stress and septic shock. *J. Microbiol. Res. Rev.* 2,11.

13. Sikka, S.C., 2001. Relative impact of oxidative stress on male reproductive function. *Curr. Med. Chem.* 8, 851-862.
14. Ochsendorf, F.R., 1999. Infections in the male genital tract and reactive oxygen species. *Hum. Reprod. Update.* 5, 399-420.
15. Li, Z., Zhang, D., He, Y., Ding, Z., Mao, F., et al., 2016. Lipopolysaccharide Compromises Human Sperm Function by Reducing Intracellular cAMP. *Tohoku J. Exp. Med.* 238, 105-112.
16. Abd-Allah, A.R., Helal, G.K., Al-Yahya, A.A., Aleisa, A.M., Al-Rejaie, S.S., et al., 2009. Pro-inflammatory and oxidative stress pathways which compromise sperm motility and survival may be altered by L-carnitine. *Oxid. Med. Cell Longev.* 2, 73-81.
17. Nezić, L., Skrbić, R., Dobrić, S., Stojiljković, M.P., Satara, S.S., et al., 2009. Effect of simvastatin on proinflammatory cytokines production during lipopolysaccharide-induced inflammation in rats. *Gen. Physiol. Biophys.* 28, 119-126.
18. Fernandez-botran, R., Gorantala, V., Sun, X., Ren, X., Perez-Abadia, G. et al. 2002. Targeting of glycosaminoglycan-cytokine interactions as a novel therapeutic approach in allotransplantation. *Transplantation* 74, 623-629.
19. Beutler, E., Duron, O. and Kelly, B., 1963. Improved method for the determination of blood glutathione. *J. Lab. Clin. Med.* 61, 882-888.
20. Habig, W.H., Pabst, M.J. and Jakoby, W.B., 1976. Glutathione S-transferase AA from rat liver. *Arch. Biochem. Biophys.* 175, 710-716.
21. Nishikimi, M., Rao, N.A. and Yagi, K., 1972. The occurrence of superoxide anion in the reaction of reduced phenazine methosulfate and molecular oxygen. *Biochem. Biophys. Res. Commun.* 46, 849-854.
22. Koracevic, D., Koracevic, G., Djordjevic, V., Andrejevic, S., Cosic, V., et al., 2001. Method for the measurement of antioxidant activity in human fluids. *J. Clin. Pathol.* 54, 356-361.
23. Aebi, H., 1984. Catalase in vitro. *Methods Enzymol.* 105, 121-126.
24. Draper, H., Hadley, M., 1990. Malondialdehyde determination as index of lipid peroxidation. *Methods Enzymol.* 186, 421-431.
25. Montgomery, H., Dymock, J.F., 1961. The determination of nitrate in water. *Analyst.* 86, 414-416.
26. Tribukait, B., 1987. Flow cytometry in assessing the clinical aggressiveness of genito-urinary neoplasms. *World J. Urol.* 5, 108-122.
27. Bancroft, J. D., Stevens, A., 1996. Theory and practice of histological technique. 4th Ed. Churchill, Livingston, Edinburgh, London, Melbourne, NY. .
28. Kumagai, A., Kodama, H., Kumagai, J., Fukuda, J., Kawamura, K., et al., 2002. Xanthine oxidase inhibitors suppress testicular germ cell apoptosis induced by experimental cryptorchidism. *Mol. Hum. Reprod.* 8, 118-123.
29. Aitken, R.J., Roman, S.D., 2008. Antioxidant systems and oxidative stress in the testes. *Adv. Exp. Med. Biol.* 636, 154-71.
30. Hales, D.B., Allen, J.A., Shankara, T., Janus, P., Buck, S., et al., 2005. Mitochondrial function in Leydig cell steroidogenesis. *Ann. N Y. Acad. Sci.* 1061, 120-134.
31. Naughton, C.K., Nangia, A.K., Agarwal, A., 2001. Pathophysiology of varicoceles in male infertility. *Hum. Reprod. Update* 7, 473-481.
32. Agarwal, A., Gupta, S., Sikka, S., 2006. The role of free radicals and antioxidants in reproduction. *Curr. Opin. Obstet. Gynecol.* 18, 325-332.
33. Rao, F., Tian, H., Li, W., Hung, H., Sun, F., et al., 2016. Potential role of punicalagin against oxidative stress induced testicular damage. *Asian J. Androl.* 18, 627-32.
34. Zhao, L., Chen, Y.-H., Wang, H., Ji, Y.L., Ning, H., et al., 2008. Reactive oxygen species contribute to lipopolysaccharide-induced teratogenesis in mice. *Toxicol. Sci.* 103, 149-157.
35. Jonsson, C.K., Setchell, B.P., Martinelle, N., Svechnikov, K., Söder, O., et al., 2001. Endotoxin-induced interleukin 1 expression in testicular macrophages is accompanied by downregulation of the constitutive expression in Sertoli cells. *Cytokine* 14, 283-288.

36. Yamada, H., Arai, T., Endo, N., Yamashita, K., Fukuda, K., et al., 2006. LPS-induced ROS generation and changes in glutathione level and their relation to the maturation of human monocyte-derived dendritic cells. *Life Sci.* 78, 926-933.
37. Jaja-Chimedza, A., Gantar, M., Mayer, G.D., Gibbs, P.D and Berry, J.P., 2012. Effects of cyanobacterial lipopolysaccharides from microcystis on glutathione-based detoxification pathways in the zebrafish (*Danio rerio*) embryo. *Toxins* 4, 390-404.
38. Sheik Abdulazeez, S., Thiruvengadam, D., 2013. Effect of lycopene on oxidative stress induced during D-galactosamine/lipopolysaccharide-sensitized liver injury in rats. *Pharm. Biol.* 51, 1592-1599.
39. Skibska, B., Józefowicz-Okonkwo, G., Goraca, A., 2006. Protective effects of early administration of alpha-lipoic acid against lipopolysaccharide-induced plasma lipid peroxidation. *Pharmacol Rep.* 58, 399-404.
40. Spolarics, Z., Wu, J-X., 1997. Role of glutathione and catalase in H₂O₂ detoxification in LPS-activated hepatic endothelial and Kupffer cells. *Am. J. Physiol.* 273, G1304-G1311.
41. Reddy, M.M., Mahipal, S.V., Subhashini, J., Reddy, M.C., Roy, K.R. et al., 2006. Bacterial lipopolysaccharide-induced oxidative stress in the impairment of steroidogenesis and spermatogenesis in rats. *Reprod. Toxicol.* 22, 493-500.
42. Pigeolet, E., Corbisier, P., Houbion, A., Lambert, D., Michiels, C., et al., 1990. Glutathione peroxidase, superoxide dismutase, and catalase inactivation by peroxides and oxygen derived free radicals. *Mech. Ageing Dev.* 51, 283-97.
43. Wu, Q.J., Wang, Y.Q., Qi, Y.X., 2016. The protective effect of procyanidin against LPS-induced acute gut injury by the regulations of oxidative state. *Springerplus.* 5, 1645.
44. Li, L., Ma, P., Liu, Y., Huang, C., O, W.S., et al., 2013a. Intermedin attenuates LPS-induced inflammation in the rat testis. *PLoS One* 8, e65278.
45. Requintina, P.J., Oxenkrug, G.F., 2003. Differential effects of lipopolysaccharide on lipid peroxidation in F344N, SHR rats and BALB/c mice, and protection of melatonin and NAS against its toxicity. *Ann. N Y. Acad. Sci.* 993, 325-333.
46. Allen, J.A., Diemer, T., Janus, P., Hales, K.H. and Hales, D.B., 2004. Bacterial endotoxin lipopolysaccharide and reactive oxygen species inhibit Leydig cell steroidogenesis via perturbation of mitochondria. *Endocrine.* 25, 265-275.
47. Aly, H., Lightfoot, D.A. and El-Shemy, H.A., 2009. Modulatory role of lipoic acid on lipopolysaccharide-induced oxidative stress in adult rat Sertoli cells in vitro. *Chem Biol Interact.* 182, 112-118.
48. Hu, W., Shi, L., Li, M.Y., Zhou, P.H., Qiu, B., et al., 2017. Adrenomedullin protects Leydig cells against lipopolysaccharide-induced oxidative stress and inflammatory reaction via MAPK/NF- κ B signalling pathways. *Sci Rep.* 7, 16479.
49. Chang, C-Y., Tucci, M., Baker, R.C., 2000. Lipopolysaccharide-stimulated nitric oxide production and inhibition of cell proliferation is antagonized by ethanol in a clonal macrophage cell line. *Alcohol* 20, 37-43.
50. O'Bryan, M.K., Schlatt, S., Gerdprasert, O., Phillips, D.J., de Kretser, D.M., et al., 2000. Inducible nitric oxide synthase in the rat testis: evidence for potential roles in both normal function and inflammation-mediated infertility. *Biol. Reprod.* 63, 1285-1293.
51. Stangel, M., Zettl, U.K., Mix, E., Zielasek, J., Toyka, K.V., et al., 1996. H₂O₂ and nitric oxide-mediated oxidative stress induce apoptosis in rat skeletal muscle myoblasts. *J. Neuropathol. Exp. Neurol.* 55, 36-43.
52. Print, C.G., Loveland, K.L., 2000. Germ cell suicide: new insights into apoptosis during spermatogenesis. *Bioessays.* 22, 423-430.
53. Li, X., Fang, P., Mai, J., Choi, E.T., Wang, H., et al., 2013b. Targeting mitochondrial reactive oxygen species as novel therapy for inflammatory diseases and cancers. *J. Hematol. Oncol.* 6, 19.
54. Oyinloye, B.E., Adenowo, A.F., Kappo, A.P., 2015. Reactive oxygen species, apoptosis, antimicrobial peptides and human inflammatory diseases. *Pharmaceuticals (Basel)* 8, 151-175.

55. Baud, V., Karin, M., 2001. Signal transduction by tumor necrosis factor and its relatives. *Trends in Cell Biol.* 11(9), 372-7.56.
56. Hamada, E., Nishida, T., Uchiyama, Y., Nakamura, J., Isahara, K. et al., 1999. Activation of Kupffer cells and caspase-3 involved in rat hepatocyte apoptosis induced by endotoxin. *J. Hepatol.* 30, 807-818.
57. Kajihara, T., Okagaki, R. and Ishihara, O., 2006. LPS-induced transient testicular dysfunction accompanied by apoptosis of testicular germ cells in mice. *Med. Mol. Morphol.* 39, 203-208.
58. Zhou, P.H., Hu, W., Zhang, X.B., Wang, W., Zhang, L.J., 2016. Protective Effect of Adrenomedullin on Rat Leydig Cells from Lipopolysaccharide-Induced Inflammation and Apoptosis via the PI3K/Akt Signaling Pathway ADM on Rat Leydig Cells from Inflammation and Apoptosis. *Mediators Inflamm.* Article ID 7201549, 16 pages. <http://dx.doi.org/10.1155/2016/7201549>.
59. Zhan, X.X., Hu, Y.A., Qing, X.R., Mo, D.S., Cai, H.C. et al., 2014. Lipopolysaccharide affects testicular histology and reproductive endocrine function in male rats. *Zhonghua Nan Ke Xue* 20, 304-308.
60. Bosmann, H.B., Hales, K.H., Li, X., Stocco, D.M., Hales, D.B., 1996. Acute in vivo inhibition of testosterone by endotoxin parallels loss of steroidogenic acute regulatory (StAR) protein in Leydig cells. *Endocrinol.* 137,