

Raw pH fall-out as a sign of a mycorrhizal modifier of *Sorghum sudanensis*

Giorgio Masoero ^{1,*}, Pier Giorgio Peiretti ², Alberto Cugnetto ¹, Giusto Giovannetti ³

¹Accademia di Agricoltura di Torino; Torino, Italy.

²Istituto di Scienze delle Produzioni Alimentari, C.N.R., Grugliasco, Italy.

³Centro Colture Sperimentali, CCS-Aosta S.r.l., Quart, Italy.

Abstract

The management of symbiotic Microbial Biota (MB) in the soil as agents that promote the yield and health of crops, is aimed at inducing modifications of the phenotype of plants, both over and under the ground. It is here shown, in *Sorghum sudanensis* plants, that: i) a simple response to MB inoculation is the result of the fall out of the raw pH; ii) the simple NIR scans of leaves can be considered to rapidly classify the outcomes; iii) the raw pH can be considered a key-variable of leaf modifications. An experiment was carried out on *Sorghum sudanensis*. The plants were seeded in pots and grown for 66 d, and then a control non-inoculated group (C) was compared with thirteen *Arbuscular Mycorrhizae* (AM) *Glomus* inoculated groups and with two commercial MB products. A total of 374 raw pH measurements conducted on the leaves showed that the 5.18 pH units in the C group were scaled by -1.9% ($P < 0.0336$) in the MB group and by -3.4% in the AM group ($P < 0.0001$), with a relevant diversity between groups. Direct discrimination of these three groups, by means of smart NIR-SCIO, showed a % reclassification of the C, MB and AM groups of 74%, 59% and 96% in the fresh leaves and of 65%, 51% and 94% in the dried ground leaves, respectively. The composition of the dried leaves, based on a set of 14 variables predicted via NIRS models, plus the total foliar dry weight and percentage, showed a typical increase in protein, ash and hemicellulose, and a typical decrease in the cellulose, dry matter, crude fiber and crop maturity index. These variables were related to the foliar pH, as a key-variable, by means of a PLS standard model (R^2 0.81) in which a low pH steadily favored the dry mass weight and, to a lesser extent, the hemicellulose and the digestible NDF contents; on the other hand, a high pH increased the dry matter percentage and the cellulose content of the leaves. As expected, the leaves of the inoculated plants showed a more juvenile ontogenic status. The epigeal botanical modifications can be considered harmonic expressions of a luxuriant symbiosis, as testified by the homologous NIR categorization. The outlook for a symbiotic agriculture, with mycorrhizal plants, should consider the raw pH as a multifaceted variable.

Corresponding author: Giorgio Masoero, Accademia di Agricoltura di Torino, Via A. Doria 10, 10123, Torino, Italy.

Running Title: Raw pH is a signal of a mycorrhizal modifier

Key words: raw pH, arbuscular mycorrhizal fungi, microbial biota, NIR-Tomoscopy, smart NIR-SCIO, NIRS foliar discrimination, leaf composition, symbiotic agriculture.

Received: Jul 30, 2018

Accepted: Aug 13, 2018

Published: Aug 15, 2018

Editor: Prem Narain, Biostatistics, Bioinformatics, Genetics Bachelor of Science (Hons), Master of Science, Doctor of Philosophy, Doctor of Science, USA.

Introduction

A credible increase of 50% in the world feed and food necessities can be expected at the end of the next thirty years¹, and more fertilizers, water and pesticides will be needed to satisfy such needs. Starting all over again with a "Symbiotic agriculture" is a seductive "manifesto" for both a sustainable development of arable lands and for the new off-land agrotechnology². *Arbuscular Mycorrhizae* (AM) need to be closely involved in a road map of sustainable agriculture as a result of their symbiotic capabilities in the agrarian multiverse, characterised by these highlights: 1) soil preservation, with restoration of the organic matter in both topsoil and polluted soils³; 2) anticipation of the *Mitscherlich* curve of phosphorus efficiency⁴; 3) resolution of the "nitrogen paradox" as a result of the uptake of organic-N⁵; 4) benefits for the atmosphere and for the foreseen climate changes, with an important reduction in N₂O emissions⁶; 5) luxuriance of the epigeal as well as of the hypogean plant systems⁷; 6) enhancements of the functional properties of feeds and foods⁸⁻¹²; 7) induction of greater intrinsic resistance to the onset of a pathological state, regardless of the origin and nature of the plants¹³.

Experiments with AM may be divided into two categories, that is, practical and scientific⁷. The first type analyses pure soils and substrates, such as those in a microbial desert, without agrarian biotas, while the second one studies extensive and multiple interactions in the presence of immense microbial charges¹⁴.

In a first field experiment with maize fertilized by a Micosat F® complex microbial consortium¹⁵, it was found that the *in vivo* raw pH of a plant was acidified according to a de-gradient from the roots (pH -7% in mycorrhizal maize) to the stem at an ear height (-4%). Other researches in vineyards confirmed the tendency of the raw pH¹⁶, measured at the petiole insert, to be lowered after microbial fertilisation with 10 kg ha⁻¹ of MB compound inoculated near the secondary roots. The authenticity of the raw pH parameter was highlighted by Cornelissen et al.¹⁷, who considered the leaf pH as a new plant trait that could explain variations in the foliar chemistry and carbon cycling processes of subarctic plant species and types. The first work was followed by a microcosm experiment¹⁸, the results of which upset the myth of a pH dominance of the soil: the pH in fresh

leaves will only grow for +0.036 pH_{soil}⁻¹ (from -0.5 to -0.7%).

The aim of the present study has been to ascertain whether the decrease in the varied raw pH in leaves is due to AM or to a whole complex of a microbial consortium, and to assess how AM can be considered effective modifier agents of the *Sorghum sudanensis* phenotype, while considering the raw pH as a key-variable.

Experimental Procedure

Two rapid methods: the measurement of the raw pH, according to Masoero and Giovannetti¹⁵, and the smart NIR-SCIO measurement of the leaf, according to Masoero et al.¹⁹, have been used in this study to analyse the multifaceted effects of an inoculation of *Sorghum sudanensis* plants with thirteen AM *Glomus* strains and two microbial biotas fertilisers (MB), and the results have been compared with a non-inoculated control (C). Basically, we considered the *in-vivo* raw pH as a rapid indicator of symbiosis. In parallel, we considered the NIR spectra, reflected from the fresh and dried leaves (NIR-Tomoscopy), as a categorically discriminant tool. The third step concerned several traits of the leaf composition, which were predicted from an extended NIR-IR spectrum of the dried and ground tissues. Finally, the raw pH was connected, as a key-variable, to the phenotypes of plants by means of linear model analyses.

Materials and Methods

Plant Material, Inocula and Growth Conditions

Three categories were considered: i) *Arbuscular Mycorrhizae* (AM), with thirteen theses of crude inocula belonging to the *Glomus* genre from the USA (AM01-AM13); ii) Microbial Biota of the soil (MB), with two commercial approved bio-fertilizer consortia (MB1-MB2); iii) a non-inoculated Control thesis (C) (Tab. 1).

Sorghum sudanensis seeds from cv "Puma" (Padana Sementi Elette S.r.l. Tombolo, Italy) were inoculated at seeding in 10 L pots (three seeds per pot), using 12 g pot⁻¹ of crude inoculum. The growth substrate was volcanic lapillus rock (5 L) and peat (5 L). The substrate was kept in the greenhouse and drip irrigation was adopted.

Table 1. Setup of the Categories and Theses.

Categories	Theses	Species
AM - Arbuscular Mycorrhizae	AM01	<i>Claroideoglopus etunicatum</i> YV PA137
AM - Arbuscular Mycorrhizae	AM02	<i>Paraglopus occultum</i> CL700A
AM - Arbuscular Mycorrhizae	AM03	<i>Acaulospora morrowiae</i> CL290
AM - Arbuscular Mycorrhizae	AM04	<i>Gigaspora gigantea</i> PA125
AM - Arbuscular Mycorrhizae	AM05	<i>Sclerocystis sinuosa</i> MD126
AM - Arbuscular Mycorrhizae	AM06	<i>Septoglopus constrictum</i> FL328
AM - Arbuscular Mycorrhizae	AM07	<i>Gigaspora rosea</i> NY328A
AM - Arbuscular Mycorrhizae	AM08	<i>Funneliformis mosseae</i> IT201
AM - Arbuscular Mycorrhizae	AM09	<i>Rhizophagus intraradices</i> CA502
AM - Arbuscular Mycorrhizae	AM10	<i>Archaeospora trappei</i> FL208
AM - Arbuscular Mycorrhizae	AM11	<i>Scutellospora pellucida</i> MN408A
AM - Arbuscular Mycorrhizae	AM12	<i>Claroideoglopus claoideum</i> ON393
AM - Arbuscular Mycorrhizae	AM13	<i>Diversispora eburnea</i> AZ420
MB - Microbial Biota [§]	MB1	Micosat F© (CCS-Aosta, Quart, Italy)
MB - Microbial Biota [£]	MB2	MycUp© (Symborg, Murcia, Spain)
C- Control not inoculated	C	

[§]MB from finely ground cultivated *Sorghum sudanensis* roots, containing spores and *ifae* of *Funneliformis coronatus* GO01 and GU53, *F. caledonium* GM24, *F. intraradices* GB67 and GG32, *F. mosseae* GP11 and GC11, *F. viscosum* GC41; saprotrophic fungi: *Streptomyces spp.* ST60, *Streptomyces spp.* SB14, *Streptomyces spp.* SA51, *Beauveria spp.* BB48, *Trichoderma viride*, *Trichoderma harzianum* TH01, *Trichoderma atroviride* TA28, *Trichoderma spp.*; rhizosphere bacteria: *Bacillus subtilis* BA41, *Pseudomonas fluorescens* PN53, *Pseudomonas spp.* PT65 and *Pochonia chlamidosporia*, in the relative percentage of 40% crude inoculum and 21.6% bacteria and saprotrophic fungi.

[£] MB from finely ground cultivated *Sorghum sudanensis* roots, containing *Glomus iranicum var. tenuihypharum* (120 propagules g⁻¹ - 1%); rhizosphere bacteria (103 UFC g⁻¹).

The leaves were harvested after 66 d and measured to establish the raw pH of the leaves by means of direct contact of the split central vein with a combined plastic-glass Double Pore Knick® electrode (Hamilton, Reno, USA) using an "XSpH 70" pH meter (Giorgio Bormac S.r.l., Carpi, Italy).

The whole leaves from each of the 16 theses were chopped and dried at 60 °C to a constant weight, air-equilibrated, ground in a Cyclotec mill and stored for later analysis by means of NIRS.

Vibrational Spectroscopy and Chemometrics

The ground leaves were repeatedly examined (three scans) with a Spectrum IdentiCheck™ FT-NIR/MIR system (Perkin-Elmer, Beaconsfield, Buck, England) and the chemical composition was predicted via NIRS, using equations that were established on twelve different forage crops, as reported in Tassone et al.²⁰.

NIR-Tomography

A smart NIR miniaturized web-based wireless spectrophotometer, SCIO version 1.2 (Consumer Physics, Tel Aviv, Israel), with a 740-1070 nm range, was used to scan the fresh leaves on the upper foliar tinge (one scan) and the ground dry matrices (10 replicates).

Statistical Analyses

An ANOVA one-way model was used to fit the effect of the categories or of the 16 theses using PROC GLM from SAS-STAT 9.0 software (SAS Institute, Inc., Cary, NC, USA), while the Tukey test was used to adjust for LSMEANS PDIF. Univariate analyses were conducted on the values of the raw pH and on the predicted composition.

The means values of the 14 variables pertaining to the 16 theses plus the total foliar dry weight and percentage, were then related to the foliar pH, as a key-variable, using a standard Partial Least Square (PLS) model (StatBox 6.5 v., Grimmer Logiciels, Paris, France) which allows two latent variables to be achieved; in this way, it was possible to identify the variables that were favoured or contrasted by the raw foliar pH variations among the theses.

A categorical discrimination of the three main groups of the fresh and dried leaves was performed by means of chemometrics of the 331-point spectra using

the SCIO Lab proprietary software, based on AKA (Also Known As), the confusion matrix, without any mathematical pre-treatment of the spectra.

Results

Raw pH

The raw pH of the leaves appeared as a very narrow-distributed variable, which was characterized by a variation coefficient of 3.8% (Tab. 2).

The normal mean value of 5.18 ± 0.03 in the leaves from the C non-inoculated plants was significantly decreased to 5.10 ± 0.02 in the MB category (-1.5%, $P=0.0336$), and decreased to a value of 5.00 ± 0.01 in the AM category (-3.5%, $P<0.001$). Moreover, the AM vs. MB difference was highly significant, with the MBs being less acidifying. When the individual theses (Tab. 3a and 3b; Fig. 1) were considered, the differences appeared more scaled.

As far as the biofertilizers are concerned, MB1 (-1.1%) was more acidifying than MB2 (-2.0%), and the AM were scaled according to three gradients of acidity: the most acidic strains, that is, below -4%, were the AM01, AM13, AM05, AM02 and AM08 strains.

NIR-Tomography, Foliar Weight and Composition

The weight of the leaves harvested from the plants was much more reduced in the non-inoculated Control than in all other theses (Tab. 3a). Moreover, the foliar composition was affected by the category of the inoculant, as calculated from Tables 3a and 3b. The MB category scored 7 significant increments and 3 decrements vs. the C, while the AM category scored 5 and 1 variations, respectively.

As can be observed in Fig. 2, the constituents augmented in the leaves of the inoculated *Sorghum sudanensis* were: crude protein, ash, digestible NDF, hemicellulose and the ether extract; the constituents reduced in contents and properties, namely dry matter, crude fiber, cellulose and crop maturity index, were increased less.

Foliar pH as the Key-Variable

The PLS standard model (Fig. 3), with $R^2=0.81$, showed that a low pH steadily favoured the dry mass weight and, to a lesser extent, the hemicellulose, digestible NDF and ether extract contents; on the other hand, a high pH tendentially increased the dry matter




Figure 1. Histogram of the % decrease in the raw pH of the inoculated theses from the non-inoculated Control, ordered by values.




Figure 2. Histogram of the % deviation in the foliar composition of the AM and MB categories from the non-inoculated control, ordered according to the AM values.

Table 2. Raw pH values (No. 374) in the three Categories.

Statistics	Categories			Prob.		
	C	MB	AM	C vs. MB	C vs. AM	MB vs. AM
LSMeans	5.18	5.10	5.00	0.0336	<.0001	0.0002
SE ±	0.03	0.02	0.01			
RMSE		0.19				
R ²		0.12				




Figure 3. Histogram of the normalized Partial Least Squares regression coefficient of the raw foliar PH




Figure 4. Histogram of the % deviation in the grass composition of the treated barley, according to Claps et al.²⁶, with MB1 from the non-inoculated control, ordered according to the values.

Table 3a. Results of the quality of *Sorghum sudanensis* leaves (No.48) [£].

Thesis	pH	Dry Mass g	Hemi cellulose %	NDF digestible %	Ether Extr. %	ADF, %	Crop Maturity Index	Crude Fiber %	Nitrogen Free Extract %
AM – Arbuscular Mycorrhizae									
AM01	4.93b	21.1	7.9	29.1	1.4	46.0	2.3	24.2	47.0
AM02	4.96b	14.3	7.1	28.9	1.7	45.5	2.1	24.6	42.5
AM03	4.97b	14.3	8.2	27.0	1.4	43.7	2.0	25.0	45.9
AM04	5.03ab	9.0	8.8	27.9	1.5	48.3	2.1	24.3	45.4
AM05	4.95b	16.0	6.1	27.4	1.7	48.8	2.4	26.5	42.7
AM06	5.08ab	17.2	6.4	27.3	1.5	45.8	2.1	24.0	45.6
AM07	5.04ab	16.9	4.8	27.7	1.4	46.9	2.2	23.9	43.9
AM08	4.97b	17.6	5.5	29.2	1.5	46.8	2.0	23.7	43.5
AM09	5.03ab	15.7	8.3	27.4	1.6	44.6	1.9	20.3	46.9
AM10	5.04ab	8.8	5.6	30.1	1.7	43.2	1.5	22.2	45.7
AM11	5.01b	15.5	5.3	29.1	1.5	47.0	2.0	23.7	43.3
AM12	5.06ab	15.5	5.4	27.7	1.5	47.4	2.3	26.4	43.3
AM13	4.94b	18.4	7.9	27.1	1.6	47.9	2.2	25.9	43.7
MB– Microbial Biota									
MB1	5.08a	12.1	7.4	28.4	1.4	42.6	1.9	20.2	46.3
MB2	5.12a	14.7	8.4	28.3	1.4	44.1	1.9	21.1	46.7
C– Control									
C	5.18a	6.7	5.6	25.0	1.3	45.5	2.3	25.8	43.9
MB/C	-1%	99%	42%	13%	10%	-5%	-18%	-20%	6%
Prob	0.03	-	0.01	0.00	0.37	0.24	0.01	0.00	0.019
AM/C%	-3%	129%	20%	12%	18%	2%	-8%	-6%	1%
Prob	<.0001	-	0.20	<.00	0.00	0.94	0.13	0.34	0.80

[£] No. Dry Matter 16; a, b, c Means followed by the same letter are not significantly different, after Tukey's HSD test at P < 0.05.

Table 3b. Quality results on *Sorghum sudanensis* leaves (No. 48).

Thesis	Ash %	<i>In vitro</i> Digestibility %	NDF non dig. %	NDF digestibility %	ADL %	NDF %	Crude Protein %	Cellulose %	Dry Matter % [£]
AM – Arbuscular Mycorrhizae									
AM01	8.0	70.4	29.6	42.3	6.3	54.5	6.7	22.1	32.6
AM14	8.7	72.0	28.0	47.3	8.8	50.9	9.1	19.9	30.3
AM06	7.6	71.1	28.9	45.6	8.8	49.6	8.7	18.2	28.9
AM02	10.1	72.5	27.5	52.3	8.4	51.1	10.1	17.5	29.7
AM09	9.5	71.6	28.4	44.6	8.4	51.4	9.3	19.1	28.9
AM03	9.1	71.8	28.2	43.1	7.9	51.2	9.6	22.4	29.4
AM12	9.6	71.4	28.6	44.1	8.5	51.6	9.3	18.8	29.7
AM10	10.3	72.2	27.8	44.9	7.8	52.2	11.5	24.5	29.3
AM05	7.3	72.0	28.0	46.1	8.8	54.1	9.8	22.4	29.0
AM11	10.8	76.6	23.4	52.9	8.2	50.4	9.5	19.7	28.8
AM08	8.5	71.1	28.9	40.3	7.0	52.4	8.3	20.6	31.0
AM13	7.7	69.6	30.4	43.3	9.5	52.1	8.3	21.0	33.0
AM07	8.6	70.9	29.1	43.7	8.0	53.3	9.6	22.2	30.2
MB - Microbial Biota									
MB1	10.5	72.9	27.1	49.4	7.3	49.3	10.5	23.8	29.4
MB2	10.3	72.7	27.3	47.7	6.8	53.0	10.4	22.9	30.8
C – Control non-inoculated									
C	7.1	70.0	30.0	42.3	7.5	52.7	7.1	23.6	33.6
MB/C%	46%	4%	-10%	15%	-7%	-3%	48%	-1%	-10%
Prob	0.00	0.06	0.07	0.13	0.37	0.57	0.01	0.99	-
AM/C%	25%	3%	-6%	7%	9%	-1%	30%	-12%	-11%
Prob	0.03	0.12	0.12	0.48	0.78	0.89	0.03	0.03	-

[£] No. Dry Matter 16

Table 4. Reclassification of the AKA matrices of the NIR-SCIO spectra from fresh and dry leaves into three Categories, and Probability of the diagonal elements.

Categories		NIR-Tomoscopy (No. 480)			Dry leaves (No. 48)		
		C	MB	AM	C	MB	AM
Arbuscular Mycorrhizae (AM)		22%	31%	96%	30%	48%	94%
Microbial Biota (MB)		5%	59%	2%	5%	51%	3%
Control (C)		74%	11%	1%	65%	0%	1%
Prob. Diagonal elements	P	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001

percentage and the cellulose content of the leaves.

NIR-Tomography

In general, the discriminations of the three categories when the SCIO instrument was used were highly significant (Tab. 4). The leaves from plants inoculated with AM were individuated more correctly, in the scans on the fresh leaves (96%) as well as in the scans on the dry leaves (94%); the performances pertaining to the MB leaves were significant, albeit not so high (59 and 51%, respectively), while the Control leaves were individuated more clearly (74 and 65%, respectively).

Discussion

Mycorrhizal Modifier

A luxuriating effect has been displayed in foliar development. A meta-analysis of the field studies on the responses of wheat to AM has highlighted that field AM inoculation can be proposed as an effective agronomic practice for wheat production⁷, with aboveground biomass increases of around 20%, as already assessed in maize under high^{11,15,22-24} or low²⁵ Italian input conditions and under Indian conditions²¹

As expected, a more juvenile ontogenic status emerged for the leaves of inoculated plants but graded among the inocula types.

Claps et al.²⁶, in field conditions, after the autumn sowing of barley (*Hordeum vulgare*) and clover (*Trifolium alexandrinum*) tanned at 1 kg ha⁻¹ with the MB1 of the present work, observed a slight increase in yield (not significant), but also significant variations in composition (Fig. 4), albeit only in the barley grass. As in the present work, the advantage was centred on crude protein (+18%), but in that case a strong reduction in fibrousness was found, which enhanced the digestibility coefficients of the NDF, crude protein and organic matter. On the other hand, a reduction in ash (-10%) disagreed with the increase observed in the present work.

Other AA have conducted experiments on AM at more mature stages, but hay and silages have rarely been studied. Uzun²² observed that no reductions in forage yield occurred in *Sorghum sudanensis* or maize, when the diminished supply of fertilizers was balanced with the action of *Glomus spp.* and the forage produced

using an AM treatment showed better chemical characteristics when harvested at an appropriate phenological stage.

As far as maize forage is concerned, Sibi et al.²⁷ observed that an AM fungus was significant, at a level of 5%, on water soluble carbohydrates, which increased by 14%. The Authors hypothesised that inoculation with an AM fungus increased the cytokinins, and, moving nutrients from other parts to the leaves tended to delay their senescence.

Raw pH fall-out as a Key-Variable

In our short modification experiment, low pH steadily favoured the dry mass weight and, to a lesser extent, the hemicellulose, digestible NDF and the ether extract contents of the leaves. If the raw pH is a real key-variable, the modifier direction in *Sorghum sudanensis* appears quite attractive for ruminants. Phyto-evolution studies^{17,28} of a subarctic flora suggest that tissue pH itself is controlled to a great extent for a given species, because of its direct or indirect functions in the plant; in fact, a low pH corresponds to poor digestibility and may therefore act as an anti-herbivore defence in the same subarctic flora.

The field surveys of AM communities over a wide range of soil pH suggest that it is also the major driving force in structuring these communities²⁹.

It was found, in the microcosm experiments⁶, that N₂O emissions were cumulatively 42.4% higher without AM than with AM microcosms. However, it should be recalled that the Authors, in their tomato experiment, observed an increase in soil pH after an AM inoculum of 7.75 vs. 7.62 (+1.6%; P=0.004). Unfortunately, the raw pH was not measured in that experiment. In light of our present and previous results^{15,16}, it is possible to hypothesize that the transfer of [H⁺] from the soil to the AM plants could have effectively reduced the circulating acidity of the soil. Several results support this hypothesis. Bago et al.³⁰ demonstrated, in a sterile culture with *Glomus intraradices*, that extra-radical mycelium utilised NO₃-N and increased pH of the medium by up to one unit. Li and Christie³¹ showed that soil solution Zn concentrations were lower and pH values were higher in mycorrhizal treatments than non-mycorrhizal controls, and the AM effects were more pronounced at higher Zn

application rates.

Vibrational Spectroscopy

The direct NIRS discrimination of bio-fertilized crops is rare. A first work on *Ocimum basilicum* cv. "Genovese"³² assessed that three AMs were modifiers of growth, and distributors of glandular trichomes and of the essential oil composition, with a homologous relative pattern that was appreciated by NIRS and Electronic Nose. A smart NIRS technique that has recently been applied to hay-litter-bags¹⁹ has produced promising results in discriminating the fingerprint effects of bio-fertilizers. The results of direct NIR discrimination have instead been obtained for several cases. Szuvandzsiev et al.³³ suggested that the effect of very different irrigation regimes on tomato leaves is detectable by means of NIRS, irrespective of the cultivar and phenological stage; around 960 nm emerged a good correlation between different irrigation regimes and the reflectance of the raw spectra of tomato leaves taken from the 900-1000 nm representative spectral range. Foliar UV-Vis-NIR spectra (325-1075 nm) was used to discriminate high oil and high protein from a normal maize type on five sampling dates over two years³⁴; the results showed that the qualitative models constituted a robust classification method; however, the best method on average attained only 58.3±5.5% of validation, a superior value to the hazard threshold of 33%, but of no use for traceability purposes. The somewhat optimistic values reported in the present work, obtained using the smart NIR-SCIO (Tab. 4), need to be confirmed over some years and under different conditions.

Conclusions

In a meta-analysis concerning nearly half a million species x sites worldwide on twenty-one plant traits, Moles et al.³⁵ stated that the mean annual temperature was correlated more with the plant traits than with the mean annual precipitation. Considering the present results, a further recommended easy trait that could be considered is the raw pH.

In this work, the fall out of the raw pH responded as a simple sign of AM inoculation and activities. However, various responses were observed for the same *Sorghum sudanensis* with different AMs, thus suggesting more useful combinations and interactions, where the raw pH can be considered a key-variable of

plant modifications in an AM framework.

The vibrational spectroscopy of leaves can be used to rapidly classify the outcomes linked to symbiotic agents, and a smart network can be used to capitalize on useful information in an NIRS leaf data-base.

From a practical point of view, it may be recommended to monitor raw pH for traceability purposes in an AM multiplication framework.

Pursuing the study of pH from a plant-to-soil point of view is better than the inverse, and each search conducted to connect the microbial soil fingerprint (hay-litter-bags¹⁹ or others³⁶) to the NIRS fingerprint and productivity traits of leaves can be considered useful.

Acknowledgments

The authors wish to thank the Fondazione CRT, Torino – Italy, for the financial support to the scientific activities of the "Accademia di Agricoltura di Torino".

References

1. Abhilash, PC, Tripathi, V, Edrisi, SA, Dubey, RK, Bakshi, M, Dubey, PK, Singh, HB and Ebbs, SD. (2016). Sustainability of crop production from polluted lands. *Energy, Ecology and Environment*, 1: 54-65.
2. Vurukonda, SSK, Giovanardi, D and Stefani, E. (2017). Symbiotic Agriculture: increasing knowledge on the mode of action of beneficial microorganisms. Poster at the XXIII National Conference of Italian Society of Plant Pathology. Retrieved from: https://www.researchgate.net/publication/320271396_Symbiotic_Agriculture_Plant_Growth_Promotion_and_Biocontrol_Activity_of_Beneficial_Microorganisms
3. Weyens, N, van der Lelie, D, Taghavi, S, Newman, L and Vangronsveld, J. (2009). Exploiting plant-microbe partnerships to improve biomass production and remediation. *Trends in Biotechnology*, 27: 591-598.
4. Wang, XX. (2016). Variation in phosphorus acquisition efficiency among maize varieties as related to mycorrhizal functioning. Doctoral dissertation, Wageningen University, 168 pp.
5. Thirkell, TJ, Cameron, DD and Hodge, A. (2016).

- Resolving the 'nitrogen paradox' of arbuscular mycorrhizas: fertilization with organic matter brings considerable benefits for plant nutrition and growth. *Plant, Cell & Environment*, 39: 1683-1690.
6. Bender, SF, Plantenga, F, Neftel, A, Jocher, M, Oberholzer, HR, Köhl, L, Giles, M, Daniell, TJ and Van Der Heijden, MG. (2014). Symbiotic relationships between soil fungi and plants reduce N₂O emissions from soil. *The ISME Journal*, 8: 1336-1345.
 7. Pellegrino, E, Öpik, M, Bonaria, E and Ercoli, L. (2015). Responses of wheat to arbuscular mycorrhizal fungi: A meta-analysis of field studies from 1975 to 2013. *Soil Biology and Biochemistry*, 84: 210-217.
 8. Bona, E, Lingua, G and Todeschini, V. (2016) Effect of bioinoculants on the quality of crops. In: "Bioformulations: for Sustainable Agriculture" (Eds. Arora, N, Mehnaz, S and Balestrini, R), Springer, New Delhi, 93-124.
 9. Raiola, A, Tenore, GC, Petito, R, Ciampaglia, R and Ritieni, A. (2015). Improving of nutraceutical features of many important Mediterranean vegetables by inoculation with a new commercial product. *Current Pharmaceutical Biotechnology*, 16: 738-746.
 10. Migliorini, P, Torri, L, Whittaker, A, Moschini, V, Benedettelli, S and Masoero, G. (2018). Old and new common wheat (*Triticum aestivum* L.) varieties in organic: connecting agronomic, microorganism, phytochemical and bread sensory characteristics. *Journal of Food, Agriculture and Environment*, 16: 22-27.
 11. Tripaldi, C, Novero, M, Di Giovanni, S, Chiarabaglio, PM, Lorenzoni, P, Meo Zilio, D, Palocci, G, Balconi, C and Aleandri R. (2017). Impact of Mycorrhizal Fungi and rhizosphere microorganisms on maize grain yield and chemical composition. *Pakistan Journal of Agricultural Sciences*, 54: 857-865.
 12. Peiretti, PG, Tassone, S, Masoero, G and Barbera, S. (2018). Chemical and physical properties of meat of bulls and steers fed Mycorrhizal or Conventional corn. In: "Agricultural Research Updates" (Eds: Gorawala, P and Mandhatri, S), Nova Science Publishers, Inc., NY, Volume 23: 177-196.
 13. Nuti, M and Giovannetti, G. (2015). Borderline products between bio-fertilizers/bio-effectors and plant protectants: the role of microbial consortia. *Journal of Agricultural Science and Technology A*, 5: 305-315.
 14. Malusà, E, Pinzari, F and Canfora, L. (2016). Efficacy of biofertilizers: challenges to improve crop production. In: Singh, DP, Singh, H and Prabha, R. (eds.), *Microbial Inoculants in Sustainable Agricultural Productivity*, 2: 17-40.
 15. Masoero, G and Giovannetti, G. (2015). In vivo Stem pH can testify the acidification of the maize treated by mycorrhizal and microbial consortium. *Journal of Environmental & Agricultural Sciences*, 3: 23-30;
 16. Masoero, G, Cugnetto, A and Giovannetti, G. (2017). Consorzi microbici, riduttori del pH in vivo, contro la Flavescenza Dorata: primi risultati in Piemonte. *L'Enologo*, 3: 79-82.
 17. Cornelissen, JHC, Qusted, HM, Van Logtestijn, RSP, Pérez-Harguinde-guy, N, Gwynn-Jones, D, Díaz, S, Callaghan, TV, Press, MC and Aerts, R. (2006). Foliar pH as a new plant trait: can it explain variation in foliar chemistry and carbon cycling processes among subarctic plant species and types? *Oecologia*, 147: 315-326.
 18. Cornelissen, JGC, Sibma, F, Van Logtestijn, RSP, Broekman, RA and Thompson, K. (2011). Leaf pH as a plant trait: species-driven rather than soil-driven variation. *Functional Ecology*, 25: 449-455.
 19. Masoero, G, Delmastro, M, Cugnetto, A, Giovannetti, G and Nuti, M. (2018). NIRS footprint of bio-fertilizers from hay litter-bags. *Journal of Agronomy Research*, 1: 22-33.
 20. Tassone, S, Masoero, G and Peiretti, PG. 2014. Vibrational spectroscopy to predict in vitro digestibility and the maturity index of different forage crops during the growing cycle and after freeze- or oven-drying treatment. *Animal Feed Science and Technology*, 194: 12-25.
 21. Mäder, P, Kaiser, F, Adholeya, A, Singh, R, Uppal, HS, Sharma, AK, Srivastava, R,, Sahai, V, Aragno, M, Wiemken, A, Johri BN and Fried PM. (2011). Inoculation of root microorganisms for sustainable wheat-rice and wheat-black gram rotations in India. *Soil Biology and Biochemistry*, 43: 609-619.
 22. Uzun, P. (2016). Improvement of forage yield to

- improve dairy product quality: mycorrhizal fungi application and differentiation of forage conservation methods. Doctoral tesi, University of Naples Federico II. 1-84.
23. Uzun, P, Masucci, F, Varricchio, ML, Serrapica, F, Ottaviano, L and Di Francia, A. (2017). Effects of arbuscular mycorrhizal fungi and low fertilizer supply on forage quality, milk traits and profitability. *Italian Journal of Animal Science*, 16(suppl. 1): 219-220.
24. Masoero, G, Rotolo, L, Zoccarato, I, Gasco, L, Schiavone, A, De Marco, V, Meineri, G, Borreani, G, Tabacco, E, Della Casa, G, Faeti, V, Chiarabaglio, PM, Lanzanova C, Locatelli, S and Aleandri, R. 2018. Symbiotic corn can improve yield, reduce mycotoxins, and preserve nutritive value. In: "Agricultural Research Updates" (Eds: Gorawala P. and Mandhatri S.), Nova Science Publishers, Inc., NY, Volume 24, 117-140
25. Sabia, E, Claps, S, Morone, G, Bruno, A, Sepe, L and Aleandri, R. (2015). Field inoculation of arbuscular mycorrhiza on maize (*Zea mays* L.) under low inputs: preliminary study on quantitative and qualitative aspects. *Italian Journal of Agronomy*, 10: 30-33.
26. Claps, S, Sabia, E, Sepe, L, Rufrano, L, Morone, G, Paladino, F and Fedele, V. (2013). *In vivo* digestibility of different forage species inoculated with Arbuscular mycorrhiza spp. *Italian Journal of Animal Science*, 12 (s1): 8.
27. Sibi, M, Faryabi, E and Marzban Z. (2015). Assessment of forage corn quality intercropping with green beans under influence of *Rhizobium bacteria* and arbuscular mycorrhizal fungus. *Biodiversity and Environmental Sciences*, 6: 418-424.
28. Freschet, GT, Cornelissen, JHC, van Logtestijn, RSP and Aerts, R. (2010). Evidence of the 'plant economics spectrum' in a subarctic flora. *Journal of Ecology*, 98: 362-373.
29. Dumbrell, AJ, Nelson, M, Helgason, T, Dytham, C and Fitter, AH. (2010) Relative roles of niche and neutral processes in structuring a soil microbial community. *The ISME Journal*, 4: 337-345
30. Bago, B, Vierheilig, H, Piche, Y and Azcon-Aguilar, C. (1996). Nitrate depletion and pH changes induced by the extraradical mycelium of the arbuscular mycorrhizal fungus *Glomus intraradices* grown in monoxenic culture. *New Phytologist*, 133: 273-280.
31. Li, X and Christie, P. (2001). Changes in soil solution Zn and pH and uptake of Zn by arbuscular mycorrhizal red clover in Zn-contaminated soil. *Chemosphere*, 42: 201-207.
32. Copetta, A, Lingua, G, Berta, G, Bardi, L and Masoero, G. (2006). Three arbuscular mycorrhizal fungi differently affect growth, distribution of glandular trichomes and essential oil composition, appreciated by NIR Spectroscopy and Electronic Nose, in *Ocimum basilicum var. Genovese*. *Acta Horticulturae*, 723: 151-156.
33. Szuvandzsiev, P, Pék, Z, Szántó, C and Helyes, L. (2014). Application of VIS-NIR reflectance spectra for estimating water supply effect on open field processing tomato. In: XIII International Symposium on Processing Tomato, 191-195.
34. Kahriman, F, Demirel, K, Inalpulat, M, Egesel, CO and Genc, L. (2016). Using leaf based hyperspectral models for monitoring biochemical constituents and plant phenotyping in maize. *Journal of Agricultural Science and Technology*. 18: 1705-1718.
35. Moles, AT, Perkins, SE, Laffan, SW, Flores-Moreno, H, Awasthy, M, Tindall, ML, Sack, L, Pitman, A, Kattge, J, Aarssen, LW, et al. (2014). Which is a better predictor of plant traits: temperature or precipitation? *Journal of Vegetation Science*, 25: 1167-1180.
36. Broadfoot, M. (2016). Microbes added to seeds could boost crop production. *Scientific American*. Retrieved from: <https://www.scientificamerican.com/article/microbes-added-to-seeds-could-boost-crop-production/>